

RECEIVED
CENTRAL FAX CENTER

FEB 22 2007

App. No.: 10/014,842
Art Unit: 1617**Amendments to the Claims:**

This listing of claims will replace all prior versions and listing of claims in the application.

Listing of Claims:

1. (Currently Amended) An interesterified coconut oil obtained from interesterifying coconut oil with free fatty acids obtained from hydrolysis of triglycerides of safflower oil, said interesterified coconut oil ~~comprising~~ consisting essentially of 46 mol % of ~~omega-6 polyunsaturated fatty acids~~ linoleic acid, and 17 mol % of ~~Lauric~~ lauric acid, 11% myristic acid, 9% palmitic acid, 2% stearic acid, and 15% oleic acid.
2. (Previously presented) The interesterified coconut oil of claim 1, wherein the lauric acid produces quick energy for critically ill patients.
3. (Previously presented) The interesterified coconut oil of claim 1, wherein the interesterified coconut oil is nutritionally beneficial in being hypocholesterolemic and hypotriglyceridemic.
4. (Previously presented) The interesterified coconut oil of claim 1, wherein the interesterified coconut oil reduces the total cholesterol level in serum by 10% and the total cholesterol level in liver by 36%.
5. (Previously presented) The interesterified coconut oil of claim 1, wherein the recovery of the interesterified coconut oil from the interesterification reaction is in the range of 88-92%.

App. No.: 10/014,842
Art Unit: 1617

6. (Previously presented) The interesterified coconut oil of claim 1, wherein the interesterified coconut oil comprises n-6 PUFA to increase eicosanoid production in immune compromised patients.

7. (Previously presented) The interesterified coconut oil of claim 1, wherein the interesterified coconut oil has a melting point of 12-15°C and that remains as a liquid without phase separation.

8. (Previously presented) The interesterified coconut oil of claim 1, wherein the interesterified coconut oil has safflower oil fatty acids and triaglycerols of coconut oil.

9. (Currently Amended) The interesterified coconut oil of claim 1, wherein the n-6 PUFA levels are 1.8% in the unmodified coconut oil and ~~45.5%~~ 46% in the interesterified coconut oil.

10-20. (Canceled)

21. (New) An interesterified coconut oil obtained from interesterifying coconut oil with free fatty acids obtained from hydrolysis of triglycerides of safflower oil, said interesterified coconut oil comprising 46 mol % of linoleic acid and 17 mol % of lauric acid.